

# THE GLOBAL STAR FORMATION RATE FROM THE 1.4 GHz LUMINOSITY FUNCTION

L. E. CRAM

School of Physics, University of Sydney NSW 2006, Australia; L.Cram@physics.usyd.edu.au

## ABSTRACT

The decimetric luminosity of many galaxies appears to be dominated by synchrotron emission excited by supernova explosions. Simple models suggest that the luminosity is directly proportional to the rate of supernova explosions of massive stars averaged over the past  $\approx 3 \times 10^7$  yr. The proportionality may be used together with models of the evolving 1.4 GHz luminosity function to estimate the global star formation rate density in the era  $z \lesssim 1$ . The local value is estimated to be  $0.026 \text{ M}_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}$ , some 50% larger than the value inferred from the  $\text{H}\alpha$  luminosity density. The value at  $z \approx 1$  is found to be  $0.30 \text{ M}_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}$ . The 10-fold increase in star formation rate density is consistent with the increase inferred from mm-wave, far-infrared, ultra-violet and  $\text{H}\alpha$  observations.

*Subject headings:* cosmology: observations – galaxies: evolution – radio continuum: galaxies

## 1. INTRODUCTION

Two different mechanisms are believed to energize the relativistic electrons responsible for the decimetric synchrotron emission from galaxies. In classical radio galaxies and related objects, the energy source is ultimately a non-stellar central engine, perhaps a massive black hole. In normal disk galaxies and related objects, the energy source is ultimately stellar, perhaps the totality of supernova explosions over the past  $\approx 3 \times 10^7$  yr. Although the relative importance of the two mechanisms in some galaxies is debated, it is usually not difficult to determine the dominant source in any galaxy from observations of its radio morphology, and the brightness, morphology and spectrum of any optical counterpart.

For galaxies in which the radio emission is energized by stars (hereinafter star-forming galaxies) it should be possible to infer the underlying rate of star formation from the

observed radio luminosity. Because the radio luminosity function of star-forming galaxies is well determined locally, and measurable out to  $z \approx 1$ , and because radio emission usually does not suffer absorption, the derivation of star formation rates from radio luminosities offers an important addition to other methods used to determine global star formation rates (e.g., Gallego *et al.* 1995; Rowan-Robinson *et al.* 1997; Pettini *et al.* 1997; Tresse & Maddox 1998; Treyer *et al.* 1998; Blain *et al.* 1998). On the other hand, the *quantitative* link between 1.4 GHz luminosity and the rate of star formation is rather tenuous, and further work is clearly needed to strengthen the astrophysical basis of the method.

This note derives the star formation rate density for  $z \lesssim 1$  from the observed radio luminosity function, and compares the result with other determinations of this important quantity. A Hubble constant  $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$  has been adopted, and previously published quantities dependent on  $H_0$  have been scaled if necessary.

## 2. THE STAR FORMATION RATE AT $z \approx 0$

Condon (1984a, 1984b, 1988, 1989) has determined the local radio luminosity function for the combined population of spiral and irregular galaxies in the form

$$\phi_m(L) = \log_e(10^{0.4}) \left( \frac{L}{L_0} \right)^{3/2} 10^{\left( 3.15 - \left[ \frac{\log_{10}(L) - 22.0}{1.9} \right]^2 \right)}. \quad (1)$$

Here,  $L$  is the luminosity ( $\text{W Hz}^{-1}$ ) at 1.4 GHz,  $\phi_m = \log_e(m)L\phi(L)$  is the luminosity function per unit interval in  $\log_m(L)$  and per unit volume ( $\text{Mpc}^{-3}$ ), and  $L_0 = 10^{23.54}$  is the factor converting between SI units and astronomical units. The corresponding 1.4 GHz luminosity density,  $\mathcal{L}_{1.4}$ , is given by

$$\mathcal{L}_{1.4} = \int_0^\infty L\phi(L)dL = 1.8 \times 10^{19} \text{ W Hz}^{-1} \text{ Mpc}^{-3}.$$

The uncertainty in this estimate (Condon 1989, Table 3) is a factor of  $10^{\pm 0.2}$ .

Condon and Yin (1990) and Condon (1992) have determined a relationship between the 1.4 GHz luminosity of a galaxy and its current star formation rate (SFR) which can be expressed in the form

$$SFR_{1.4} = \left( \frac{L_{1.4}}{2.35 \times 10^{22} \text{ WHz}^{-1}} \right) \text{ M}_\odot \text{ yr}^{-1}. \quad (2)$$

The conversion factor in Equation (2) corresponds to an adopted initial mass function  $\Psi(M) \sim M^{-2.35}$  with  $0.1 < M < 100 \text{ M}_\odot$ . Equations (1) and (2) imply that the local value

of the global star formation rate density is  $0.026 \text{ M}_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}$ , with an uncertainty of the order of a factor of  $10^{\pm 0.2}$ . The most luminous galaxy encountered in constructing the local luminosity function of the star-forming population has a 1.4 GHz luminosity of  $\approx 10^{24.4} \text{ W Hz}^{-1}$  (Condon 1989, Table 3); its corresponding inferred star formation rate is just over  $100 \text{ M}_\odot \text{ yr}^{-1}$ .

### 3. THE STAR FORMATION RATE FOR $z \lesssim 1$

Sub-mJy radio surveys not suffering from confusion were first made by aperture synthesis telescopes in the early 1980s. The shape of the 1.4 GHz source-count distribution revealed by these studies provided compelling evidence for significant evolution of the star-forming galaxy population. As reviewed by Condon (1988), the source count distribution normalised to the Euclidean slope [i.e., the curve  $N(S)S^{-2.5}$ ] flattens for  $S_{1.4} \lesssim 5 \text{ mJy}$ . This flattening *cannot* be interpreted in terms of the population of AGN-powered galaxies which dominate the source counts above 10 mJy, nor can it arise from a non-evolving population of star-forming galaxies having the local luminosity function presented above (Danese *et al.* 1987). The faint counts can be reconciled with the local luminosity function of star-forming galaxies by allowing significant evolution of the star-forming population in the interval  $z \lesssim 1$ . Danese *et al.* (1987; see also Rowan-Robinson *et al.* 1993; Hopkins *et al.* 1998) have shown that consistency between the local luminosity function and the faint source counts for the star-forming population can be obtained by a pure luminosity evolution model satisfying  $P_{1.4}(z) \propto (1+z)^3$ .

In studies of the entire radio source population, Condon (1984a, 1984b, 1988, 1989) has established consistency between the evolving luminosity functions of both star-forming and AGN galaxies and the radio-frequency source counts over six orders of magnitude of flux density. His model uses a density evolution factor  $g(z)$  and a luminosity evolution factor  $f(z)$  in the form

$$\phi_m(L, z) = g(z)\phi_m[L/f(z), z = 0].$$

This is the ‘co-moving’ luminosity function; Condon adopted a cosmological model with  $\Omega_0 = 1$  and  $h = 0.5$ . Values of  $f(z)$  and  $g(z)$  taken from Condon (1984b, Table 1), combined with Equations (1) and (2), lead to the  $z$ –dependence of the global star formation rate density illustrated in Figure 1. The curve is consistent with that which would be derived using pure luminosity evolution  $P(z) \propto (1+z)^3$ .

Although Condon’s (1984b) model establishes consistency between the evolving luminosity function and the source counts, it is not unique. Nevertheless, source-count models involving significant evolution of the population of sub-mJy galaxies find growing

support as studies of their optical counterparts show signs of vigorous star-formation. For example, Kron, Koo & Windhorst (1985) found that the optical counterparts of the sub-mJy population include a blue component, dubbed ‘B radio galaxies’, with peculiar optical morphologies suggestive of interactions and mergers. Benn *et al.* (1993) have shown that many faint radio sources appear to be more-distant counterparts of the IRAS star-forming galaxy population. This population probably overlaps with the ‘B radio galaxies’. *Hubble Space Telescope* images suggest that the majority of sub-mJy radio sources are starburst or post-starburst objects, embedded in luminous disk galaxies and triggered by interactions (Windhorst *et al.* 1995).

#### 4. COMPARISON WITH OTHER ESTIMATES

Gallego *et al.* (1995) have estimated the global star formation rate at  $z \approx 0$  from their observations of the local  $H\alpha$  luminosity density. Their derived star formation rate density is  $0.013 \text{ M}_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}$ , one-half the value derived above. While a factor of two uncertainty in the global star formation density derived by such different methods is not disturbingly large, it is of interest to explore the origin of the difference. Accordingly, Figure 2 compares the 1.4 GHz and  $H\alpha$  luminosity functions and the corresponding distributions of the inferred star formation rate densities, both expressed on a common  $\log(SFR)$  scale to facilitate comparison.

Over the range where the  $H\alpha$  luminosity function is well determined ( $4 \times 10^{33} < L_{H\alpha} < 4 \times 10^{35} \text{ W}$  or  $0.4 < SFR_\alpha < 40 \text{ M}_\odot \text{ yr}^{-1}$ ) the two luminosity functions cross, with the  $H\alpha$  distribution rising somewhat more steeply to lower luminosities and dropping slightly below the 1.4 GHz function at the highest observed point. At faint luminosities the extrapolation of the  $H\alpha$  luminosity function rises above the 1.4 GHz function, but the difference never exceeds a factor of 5 even at the faintest luminosities. The star formation rate density inferred from  $H\alpha$  is approximately twice the density inferred from 1.4 GHz observations, for values of SFR below about  $40 \text{ M}_\odot \text{ yr}^{-1}$ .

The excess of the 1.4 GHz value of SFR relative to the  $H\alpha$  value therefore arises at high values of SFR. This is clear from Figure 2, which shows that the  $H\alpha$  luminosity function is  $\approx 10^{-6} \text{ dex}^{-1} \text{ Mpc}^{-1}$  at  $SFR_\alpha = 100 \text{ M}_\odot \text{ yr}^{-1}$  while at the same  $SFR_{1.4}$ , the 1.4 GHz luminosity function is a factor of 25 greater. Luminosity weighting amplifies this difference in the plot of the star formation rate density. As a result, the upper quartile of star formation rate densities lies above  $25 \text{ M}_\odot \text{ yr}^{-1}$  according to the 1.4 GHz data, and above only  $15 \text{ M}_\odot \text{ yr}^{-1}$  for the  $H\alpha$  data. Even more extreme is the high star formation rate tail which extends to values  $> 100 \text{ M}_\odot \text{ yr}^{-1}$  in the 1.4 GHz data, but is absent in  $H\alpha$  (see

also Cram *et al.* 1998; objects in this category include Arp 220 and NGC 6240). The excess SFR density in the 1.4 GHz data at high values of SFR more than outweighs the excess H $\alpha$  SFR density at low values of SFR, and accounts for the somewhat higher global rate deduced from the 1.4 GHz data.

Other estimates of the global star formation rate at low redshift ( $z \lesssim 0.3$ ) have been made by Treyer *et al.* (1998) and Tresse & Maddox (1998). The former, based on balloon-borne observations at 200 nm, yields  $0.006 \text{ M}_\odot \text{ yr}^{-1}$  (adjusted to  $h = 0.5$ ) at  $z \approx 0.15$ , and the latter, based on H $\alpha$  spectra from the *I*-selected Canada-France Redshift Survey, yields  $0.020 \text{ M}_\odot \text{ yr}^{-1}$  at  $z \approx 0.2$ . The H $\alpha$  value is compatible with that of Gallego *et al.* (1995), while the far-UV value is smaller, perhaps significantly so. Both are smaller than the 1.4 GHz value.

There have been several determinations of the evolution of the SFR density beyond  $z \approx 0.25$  (e.g., Lilly *et al.* 1996; Madau *et al.* 1996; Connolly *et al.* 1997; Rowan-Robinson *et al.* 1997). All agree that there is a significant rise between  $z = 0$  and  $z = 1$ , of a factor of order 10. The 1.4 GHz results are consistent with this trend.

It should be noted that dust extinction leads to uncertainty in estimates of the star formation rate from UV and optical indicators. For example, Pettini *et al.* (1997, their Fig 7) suggest that an upward revision of a factor of approximately 2 - 6 should be applied to uncorrected estimates of star formation rates based on ultraviolet observations. Their revised values lie close to, but below, those estimated in this paper in the range  $0.2 \lesssim z \lesssim 1$ . Blain *et al.* (1998, their Figure 8) have estimated the evolution of the global star formation rate using sub-mm observations which, like radio observations, should be unaffected by extinction. Their estimates agree, to within a factor of two over the range  $z \lesssim 1$ , with those derived here.

## 5. DISCUSSION AND SUMMARY

In addition to the consistency with sub-mm observations noted above, there is close agreement between the evolution of the SFR density derived from 1.4 GHz data, and from far-infrared observations of the *Hubble Deep Field* by the Infrared Space Observatory (Rowan-Robinson *et al.* 1997). This result could have been anticipated, in view of the strong correlation between 1.4 GHz and far-infrared luminosities in local star-forming galaxies, and the viability of theories of the correlation which relate both luminosities to the number of massive stars in the galaxies (e.g. Condon 1992). Estimates of the SFR density inferred from far-UV or H $\alpha$  data (uncorrected for extinction) appear to be systematically

lower, by a factor of order 2 – 4. The discrepancy could be a consequence of a moderate amount of optical/far UV extinction, an interpretation that finds some support in studies of nearby galaxies which have revealed evidence of widespread extinction, including high attenuation in some objects (e.g., Young, Kleinmann & Allen 1988; see also Pettini *et al.* 1997). However, it should also be noted that the ratio of the instantaneous luminosity of various SFR indicators to the short-term average of the SFR itself is a complex function of time (e.g., H $\alpha$  is weighted strongly to the most massive stars’ lifetimes, while 1.4 GHz radiation is weighted to the more abundant stars lying just above the lower mass limit for supernovae). It is likely that factors other than extinction will also prove to be relevant in explaining the discrepancy between various SFR indicators.

Further studies of the extinction in galaxies having high values of  $SFR_{1.4}$  and low values of  $SFR_{\alpha}$  could help resolve this question. Of particular interest are galaxies having  $SFR_{1.4} \gtrsim 25 \text{ M}\odot \text{ yr}^{-1}$ . In such galaxies, the ‘shortfall’ in H $\alpha$  emission is particularly large, and this might lead to obvious characteristic differences between the morphology in 1.4 GHz and H $\alpha$  images.

In Condon’s (1984b) model the evolution of the star-forming galaxy luminosity function is constrained almost exclusively by the fit to sub-mJy 1.4 GHz source counts. This is not a particularly tight constraint, and as more redshifts become available for sub-mJy star-forming galaxies the evolution will become better defined. Already, the studies of Benn *et al.* (1993), Rowan-Robinson *et al.* (1993) and Hopkins (1998) suggest that consistency arises between the local 1.4 GHz luminosity function of star-forming galaxies, the sub-mJy source counts and the available  $n(z)$  distributions if there is luminosity evolution according to  $(1+z)^3$ . This picture is consistent with results obtained at other wavelengths (e.g. Hacking, Condon & Houck 1987). Since the 1.4 GHz data suggest that approximately 50% of star formation at  $z = 1$  takes place in star forming galaxies with  $L_{1.4} \gtrsim 10^{24} \text{ W Hz}^{-1}$ , and since such galaxies are not difficult to detect at radio frequencies ( $S_{1.4} \approx 250 \mu\text{Jy}$ ), optical spectroscopy of complete samples of sub-mJy 1.4 GHz sources could make an important contribution to our understanding of cosmic star formation history in the era  $z \lesssim 1$ .

It is worth noting that studies of the cosmic evolution of *powerful* radio galaxies (i.e., AGN-powered objects with  $P_{1.4\text{GHz}} > 10^{26} \text{ W Hz}^{-1}$ ) suggest that the population undergoes pure luminosity evolution according to  $P_{1.4}(z) \propto (1+z)^3$  out to  $z \approx 2$ , with a luminosity dependent redshift cut-off at higher  $z$  (e.g. Dunlop & Peacock 1990). As emphasised by Dunlop (1998), these studies provide a robust determination of the evolution of the 1.4 GHz luminosity density, despite remaining uncertainties in the high- $z$  evolution of the luminosity function itself. It is a remarkable fact that the luminosity evolution of the AGN-powered population appears to be quite similar that of the star-forming population (Danese *et*

*al.* 1987; Dunlop 1998). Dunlop (1998) advances the suggestion that the star formation rate and the AGN (black hole) fueling rate might both be related to the global rate of gravitational accretion/condensation.

The use of 1.4 GHz radiation to determine star formation rates offers some advantages over alternative methods. For example, 1.4 GHz radiation does not usually suffer absorption, and the local 1.4 GHz luminosity function of star-forming galaxies is well determined over a wide luminosity range. Compared with mm-wave and far-infrared (FIR) indicators, which share these advantages over optical and UV indicators, deep 1.4 GHz observations have superior sensitivity, and offer astrometric precision sufficient to make direct optical identifications of even the faintest sources. The main weakness is the tenuous linkage between 1.4 GHz radiation and the astrophysics of star formation. Spatially resolved observations probing the relationship between decimetric radio emission and star forming processes in selected galaxies (e.g. Duric *et al.* 1995; Marsh & Helou 1998; Smith *et al.* 1998) tend to support the model used above, but further work is clearly needed to strengthen the astrophysical basis of the method.

## REFERENCES

- Benn, C. R., Rowan-Robinson, M., McMahon, R. G., Broadhurst, T. J., & Lawrence, A. 1993, MNRAS, 263, 98
- Blain, A. W., Smail, I., Ivison, R. J. & Kneib, J.-P. 1998, MNRAS(in press)
- Condon, J. J. 1984a, ApJ, 282, 44
- Condon, J. J. 1984b, ApJ, 287, 461
- Condon, J. J. 1998, in “Galactic and Extragalactic Radio Astronomy”, eds G. L. Vershuur & K. I. Kellermann (Berlin: Springer-Verlag), 641
- Condon, J. J. 1989, ApJ, 338, 13
- Condon, J. J., & Yin, Q. F. 1990, ApJ, 357, 97
- Condon, J. J. 1992, ARA&A, 30, 575
- Connolly, A. J., Szalay, A. S., Dickinson, M., SubbaRao, M. U., & Brunner, R. J. 1997, ApJ, 486, L11
- Cram, L. E., Hopkins, A. M., Mobasher, B., & Rowan-Robinson, M. 1998, ApJ, (in press)
- Danese, L., De Zotti, G., Franceschini, A. & Toffolatti, L. 1987, ApJ, 318, L15
- Dunlop, J. S. 1998, in “Observational Astronomy with the New Radio Surveys”, eds M. N. Bremer, N. Jackson & I. Perez-Fournon (Dordrecht: Kluwer), 157

- Dunlop, J. S. & Peacock, J. A. 1990, MNRAS, 247, 19
- Duric, N., Gordon, S. M., Goss, W. M., Viallefond, F., & Lacey, C. 1995, ApJ, 445, 173
- Gallego, J., Zamorano, J., Aragón-Salamanca, A., & Rego, M. 1995, ApJ, 455, L1
- Hacking, P., Condon, J. J., & Houck, J. R. 1987, ApJ, 316, L15
- Hopkins, A. M. 1998, “The Phoenix Multiwavelength Deep Survey”, PhD Thesis, University of Sydney.
- Hopkins, A. M., Mobasher, B., Cram, L. E., & Rowan-Robinson, M. 1998, MNRAS, 296, 839
- Kron, R. G., Koo, D. C. & Windhorst, R. A. 1985, A&A, 146, 38
- Lilly, S. J., Le Fèvre, O., Hammer, F., & Crampton, D. 1996, ApJ, 460, L1
- Madau, P., Ferguson, H. C., Dickinson, M. E., Giavalisco, M., Steidel, C. C., & Fruchter, A. 1996, MNRAS, 283, 1388
- Marsh, K. A., & Helou, G. 1998 ApJ, 493, 121
- Pettini, M., Steidel, C. C., Adelberger, K. L., Kellogg, M., Dickinson, M., & Giavalisco, M. 1997, in *Origins*, ASP Conf. Ser. Vol 148, ed C. E. Woodward, J. M. Shull & H. A. Thronson, in press
- Rowan-Robinson, M., Benn, C. R., Lawrence, A., McMahon, R. G., Broadhurst, T. J. 1993, MNRAS, 263, 123
- Rowan-Robinson, M., *et al.* 1997, MNRAS, 289, 490
- Smith, H. E., Lonsdale, C. J., Lonsdale, C. J., & Diamond, P. J. 1998, ApJ, 493, L17
- Tresse, L., & Maddox, S. J. 1998, ApJ, 495, 691
- Treyer, M. A., Ellis, R. S., Milliard, B., Donas, J. & Bridges, T. J. 1998, astro-ph/9806056
- Young, J.S., Kleinmann, S.G. & Allen, L.E. 1988, ApJ, 334, L63
- Windhorst, R. A., Fomalont, E. B., Kellermann, K. I., Partridge, R. B., Richards, E., Franklin, B. E., Pascarelle, S. M. & Griffiths, R. E. 1995, Nature, 375, 471






Fig. 1.— Variation of the global star formation rate density with redshift, derived from the 1.4 GHz luminosity function of Condon (1984b).




Fig. 2.— Luminosity distributions (left panel) and star formation rate densities (right panel). Symbols denote the indicator used, and the redshift of the determination. The dash-dot curve depicts extrapolations of the local  $H_{\alpha}$  and 1.4 GHz distributions beyond the brightest and faintest detected objects. The abscissae, labelled “star formation rate”, represent a common scale for 1.4 GHz and  $H_{\alpha}$  luminosities, corresponding to  $L_{1.4}/2.35 \times 10^{22} \text{ W Hz}^{-1}$  and  $L_{H_{\alpha}}/9.4 \times 10^{33} \text{ W}$ .